

MHN

IN THE UNITED STATES DISTRICT COURT FOR
THE NORTHERN DISTRICT OF ILLINOIS
EASTERN DIVISION

FILED

FEB 05 2008 *aw*

2-5-2008

MICHAEL W. DOBBINS
CLERK, U.S. DISTRICT COURT

UNITED STATES

CASE # 82 CR 555

vs

08 CV 116

LOUIS C. SHEPTIN

1 TO: JUDGE KOROMAS
1 JUDGE KOUNLEY

URGENT
ADVISORY TO COURT

DIRECTED TO JUDGES KOROMAS & KOUNLEY:

DEFENDANT, LOUIS C. SHEPTIN FILED IN
SHEPTIN v ATTORNEY GENERAL, 08 CV 0116

AND AMENDED COMPLAINT, COMPRISED

OF A FORM; 16 ADDITIONAL PAGES; AND

THREE EXHIBITS. EXHIBIT MARKED "XX"

IS AN APPOINTMENT SLIP FROM THE

DENTAL CLINIC AT MCC CHICAGO. THE

DATE OF REQUEST FOR DENTAL CARE WAS

DECEMBER 21, 2007. APPOINTMENT DATE 1/28/08.

THE DATE SET BY THE DENTIST FOR THE
APPOINTMENT WAS JANUARY 28, 2008,

THE SAME DAY A STATUS HEARING WAS

HELD BEFORE MAGISTRATE KEYS, IN THIS CASE
DEFENDANT SHEPTIN DID NOT ATTEND FOR GOOD
CAUSE?

DEFENDANT DID NOT APPEAR AT
THE JANUARY 28, 2008 STATUS HEARING,
BECAUSE DEFENDANT HAS THREE TEETH
INFECTED, BROKEN-OFF AT GUM LINE¹
AND HAS REPEATEDLY REQUESTED DENTAL
EXTRACTION DUE TO AN ELEVATED RISK
OF CORONARY PROLAPSE/INFECTION; AND DAILY
PAIN, REPEATEDLY IGNORED! DEFENDANT DID NOT
WANT TO MISS THE APPOINTMENT.

THE DEFENDANT, HAD A CHOICE,
COURT OR DENTIST, HE CHOSE DENTIST

UNFORTUNATELY

THE DENTIST LIED⁴, PLACING
 DEFENDANT'S LIFE IN IMMINANT
 RISK, OR DANGER OF SERIOUS BODILY
 INJURY. SEE ALPHIN V. TONEY 281
 F 3d 709 (2002) DENTAL / MOUTH INFECTION
 IMMINANT DANGER. THE APPOINTMENT NEVER CALLED

THE FACT THAT DEFENDANT HAS SERIOUS
 HEART DISEASE AND HAS BEEN ANGIOPLASTY⁵
 TWICE IN THE PAST 6 WEEKS, EXPONENTIALLY
 RAISES THE DANGER TO DEFENDANT SLEEPING.

THE MEDICAL STAFF AT MCL ARE
 EITHER VERY VINDICTIVE OR SIMPLY VERY
 STUPID, IN ANY EVENT YOUR DEFENDANT'S
 JANUARY 28, 2008 APPOINTMENT AS
 DEMONSTRATED IN EXHIBIT XX ATTACHED

TO THE AMENDED COMPLAINT IN SHEPTIN
v. ATTORNEY GENERAL 08 CV 116 SPEAKS FOR
ITSELF; WAS CANCELED BY MCC!

ADD INSULT TO INJURY THE
MEDICAL STAFF TREAT DEFENDANTS DENTAL
PAIN WITH

MCC CHICAGO PHARMACY (312) 322-0567
71 W. VAN BUREN - CHICAGO, IL 60605
107401 P. HARVEY 01/31/08
SHEPTIN, LOUIS CHARLES 90355-024
MCC CHICAGO - C03-019L
TAKE ONE TABLET 3 TIMES A DAY AFTER
MEALS

IBUPROFEN 800 MG TAB
(0) Refills 01/31/2008 IL
CAUTION: Federal/State law prohibits transfer of this drug
to any person other than patient for whom prescribed.

2400 mg^{1/} OF IBUPROFEN DAILY ELEVATING
DEFENDANT'S RISK OF DEATH EVEN FURTHER!
SEE CONTRAINDICATIONS ATTACHED
AND MARKED EXHIBIT "A"

IT CLEARLY STATES ON THE ATTACHED
"PATIENT MEDICATION INFORMATION"
THAT THE RISK OF SERIOUS AND
SOMETIMES FATAL HEART PROBLEMS,
HEART ATTACK, AND STROKE MAY
BE INCREASED WITH THE USE OF
[IOPROFEN] - THIS MEDICATION IS ALSO NOT
TO BE USED CONCURRENTLY WITH BETA
BLOCKERS IE (SOTOLOL) OR ASPIRIN^{II}. IT
IS DEFINITELY CONTRAINDICATED AFTER
HEART SURGERY OF ANY TYPE INCLUDING OR
BYPASS OR ANGIOPLASTY. SEE PDR - MOTILIN P. 1125

SO, PREMISES CONSIDERED IT
IS VERY APPARENT DEFENDANT INDEED
IS AT AN ELEVATED RISK OF IMMINENT

DANGER FOR SERIOUS BODILY INJURY; DEATH
HEART ATTACK, STROKE, AND CANCERS
THE RECORDS, FILES AND EVIDENCE FILED
SHOW A PATTERN OF CONDUCT UNACCEPTABLE
IN A DEMOCRATIC SOCIETY WHERE RULE OF
LAW AND CONSTITUTION PREVAIL ALONG WITH
CIVIL RIGHTS FOR ALL; CERTAINLY CONSTITUTES ASSAULT
MAYBE WITH INTENT →
 DEFENDANT DEMANDS HIS RELEASE OR
MEDICAL TREATMENT. DEFENDANT IS AVAILING
HIMSELF FOR PHYSICAL INSPECTION OF HIS MOUTH
AS THE COURT DEMANDS JUST. PICTURE SPEAKS A THOUSAND
WORDS. . . .

DATED 2/2/08

ATTACHMENT
PATIENT INFO

RESPECTFULLY SUBMITTED,



LOUIS C. SHEPTIN
21 W. VAN BUREN ST
CHICAGO, IL 60605

PATIENT MEDICATION INFORMATION

IBUPROFEN 800 MG TAB

Rx # 107401

COMMON USES: This medicine is a nonsteroidal anti-inflammatory drug (NSAID) used to treat mild to moderate pain, osteoarthritis, and rheumatoid arthritis. It may also be used to treat other conditions as determined by your doctor.

BEFORE USING THIS MEDICINE: WARNING: THE RISK OF SERIOUS AND SOMETIMES FATAL HEART PROBLEMS, HEART ATTACK, AND STROKE may be increased with the use of this medicine. This risk may be increased the longer you use this medicine. Risk may also be higher in patients who have heart problems or who are at risk for heart problems. THIS MEDICINE SHOULD NOT BE USED to treat pain before or after coronary artery heart bypass (CABG) surgery. THE RISK OF SERIOUS AND SOMETIMES FATAL STOMACH AND BOWEL PROBLEMS, including bleeding, ulcers, and holes in the stomach and bowel, is increased while using this medicine. These problems may occur at any time during therapy, with or without symptoms. Elderly patients are at higher risk for serious stomach problems. Ask your doctor or pharmacist for more information about this medicine and its side effects. Some medicines or medical conditions may interact with this medicine. INFORM YOUR DOCTOR OR PHARMACIST of all prescription and over-the-counter medicines that you are taking. DO NOT TAKE THIS MEDICINE if you are also taking heparin or aspirin. ADDITIONAL MONITORING OF YOUR DOSE OR CONDITION may be needed if you are taking serotonin reuptake blocker medicines such as fluoxetine or citalopram, blood thinners such as warfarin, bisphosphonates such as alendronate or risedronate, cyclosporine, corticosteroids such as prednisone, high blood pressure medicines (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), water pills, diuretics such as furosemide, hydrochlorothiazide, furosemide, lithium, methotrexate, or aspirin. DO NOT START OR STOP ANY MEDICINE without doctor or pharmacist approval. Inform your doctor of any other medical conditions including poorly controlled diabetes, dehydration, heart problems (such as heart failure or history of heart attack), swelling of the hands, feet, or ankles (edema), high blood pressure, history of stroke, blood clotting problems, stomach or bowel problems (such as bleeding or ulcers), history of tobacco use or alcohol use, kidney problems, liver problems, blood or bleeding problems (such as anemia), asthma, growing in the nose (nasal polyps), any allergies (especially history of anaphylactic reaction, symptoms of low blood sugar, or food sensitivity), pregnancy, or breastfeeding. DO NOT TAKE THIS MEDICINE if you have a history of allergy to aspirin or other NSAIDs (e.g., naproxen, celecoxib). USE OF THIS MEDICINE IS NOT RECOMMENDED if you have history of severe kidney disease or if you are going to have or have recently had coronary artery heart bypass (CABG) surgery. Contact your doctor or pharmacist if you have any questions or concerns about taking this medicine.

HOW TO USE THIS MEDICINE: Follow the directions for using this medicine provided by your doctor. This medicine may come with a medication guide. Read it carefully. Ask your doctor, nurse, or pharmacist any questions that you may have about this medicine. TAKE THIS MEDICINE with a full glass (8 oz./240 ml) of water. DO NOT lie down for 30 minutes after taking this medicine. The dosage is based on your medical condition and response to therapy. If repeat doses are needed, they are usually given 6 or 8 hours apart, or as directed by your doctor. When used in children, the dose is based on your child's weight. Read the product instructions to find the appropriate dose for your child's weight. Consult the pharmacist or doctor if you have questions or if you need help in choosing the appropriate dosage form. THIS MEDICINE MAY BE TAKEN WITH FOOD if it upsets your stomach. Taking it with food may not decrease the risk of stomach or bowel problems (such as bleeding or ulcers) that may occur while taking this medicine. Ask with your doctor or pharmacist if you experience persistent stomach upset. STORE THIS MEDICINE at room temperature, away from heat and light. Do not store in the bathroom. IF YOU MISS A DOSE OF THIS MEDICINE, take it as soon as possible. If it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not take 2 doses at once.

CAUTIONS: THIS MEDICINE INCREASES YOUR RISK OF SERIOUS STOMACH OR BOWEL PROBLEMS (such as ulcers and bleeding). This risk is increased if you are elderly or are in poor health, if you have a history of smoking or drinking alcohol, if you take corticosteroid medicines (such as prednisone) or blood thinners (such as warfarin), or if you take this medicine for a long period of time. THIS MEDICINE MAY ALSO INCREASE YOUR RISK for certain serious heart and blood vessel problems (such as heart attack and stroke). TAKE THIS MEDICINE EXACTLY AS PRESCRIBED BY YOUR DOCTOR. DO NOT TAKE AT THE LOWEST POSSIBLE DOSE FOR THE LONGEST TIME PERIOD. Talk with your doctor for further information. DO NOT TAKE THIS MEDICINE IF YOU HAVE HAD A SEVERE ALLERGIC REACTION to aspirin or any medicine containing aspirin or to a nonsteroidal anti-inflammatory drug (such as feldene, Motrin, Naprosyn, Clinoril). A severe reaction includes a severe rash, hives, breathing difficulties, or dizziness. If you have a question about whether you are allergic to this medicine or if a certain medicine is a nonsteroidal anti-inflammatory drug, contact your doctor or pharmacist. DO NOT EXCEED THE RECOMMENDED DOSE or take this medicine for longer than 10 days for pain or 3 days for fever, unless directed by your doctor, pharmacist, or other medical professional. During blood counts, liver function tests, and kidney function tests, may be performed to monitor your progress or to check for side effects, especially if you are taking this medicine for a long period of time. KEEP ALL DOCTOR AND LABORATORY APPOINTMENTS while you are taking this medicine. DO NOT DRIVE, OPERATE MACHINERY, OR DO ANYTHING ELSE THAT COULD BE DANGEROUS until you know how you react to this medicine. ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take this medicine or other pain relievers/fever reducers. BEFORE YOU BEGIN TAKING ANY NEW MEDICINE, either prescription or over-the-counter, check with your doctor or pharmacist. If you are also taking aspirin, as prescribed by your doctor for reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), continue to take the aspirin and consult your doctor or pharmacist before using this medicine. CAUTION IS ADVISED WHEN USING THIS MEDICINE IN THE ELDERLY because they may be more sensitive to the effects of this medicine, especially its effects on the stomach or bowel effects (such as bleeding or ulcers), or kidney effects. FOR WOMEN: USE OF THIS MEDICINE DURING PREGNANCY has resulted in fetal and newborn death. If you think you may be pregnant, contact your doctor immediately. THIS MEDICINE IS EXCRETED IN BREAST MILK. IF YOU ARE OR WILL BE BREAST-FEEDING while you are using this medicine, check with your doctor or pharmacist to discuss the risks to your baby.

POSSIBLE SIDE EFFECTS: SIDE EFFECTS that may go away during treatment include nausea, vomiting, diarrhea, gas, constipation, indigestion, dizziness, joint pain, drowsiness, or headache. If you get a rash, hives, or are bothered by any of these, check with your doctor. CHECK WITH YOUR DOCTOR AS SOON AS POSSIBLE if you experience ringing in ears. CONTACT YOUR DOCTOR IMMEDIATELY if you experience rapid or pounding heartbeat, easy bruising or bleeding, very stiff neck, or mental/mood changes. CONTACT YOUR DOCTOR IMMEDIATELY if you experience sharp or crushing chest pain, sudden shortness of breath, sudden leg pain, sudden severe headache, vomiting, dizziness, or feeling changes in vision, numbness of an arm or leg, slurred speech, one-sided weakness, sudden unexplained weight gain, change in amount of urine produced, severe or persistent stomach pain, vomit that looks like coffee grounds, black tarry stools, itching, redness, swollen, blistered, painful, or peeling skin, yellowing of the skin or eyes, dark urine, right-sided tenderness, severe or persistent tiredness, fever, chills, or sore throat, severe or persistent nausea, swelling of hands, ankles, feet, face, lips, eyes, throat, or tongue, difficulty swallowing or breathing, or hoarseness. AN ALLERGIC REACTION TO THIS MEDICINE is unlikely, but seek immediate medical attention if it occurs. Symptoms of an allergic reaction include rash, itching, swelling, severe dizziness, or trouble breathing. If you notice other effects not listed above, contact your doctor, nurse, or pharmacist.

OVERDOSE: If overdose is suspected, contact your local poison control center or emergency room immediately. Symptoms of overdose may include severe stomach pain, coffee ground-like vomit, unusually fast or slow heartbeat, trouble breathing, extreme drowsiness, loss of consciousness, and seizures.

ADDITIONAL INFORMATION: DO NOT SHARE THIS MEDICINE with others for whom it was not prescribed. DO NOT USE THIS MEDICINE for other health conditions. KEEP THIS MEDICINE out of the reach of children and pets. IF USING THIS MEDICINE FOR AN EXTENDED PERIOD OF TIME, obtain refills before your supply runs out.

Database Edition 06.1 - Expires April 2008
Copyright 2008 Wolters Kluwer Health, Inc.
All rights reserved.

EXHIBIT "A"

BOP Rx LABEL

TRAK# WFP03